

*A review of*  
**Foundations of the  $\text{AdS}_5 \times S^5$  Superstring**

**Alexander Farren**

*supervised by* Prof. Sergey Frolov

March 12, 2024

# Why $\text{AdS}_5 \times S^5$ ?


- ▶ Modern TP: *Can we reconcile Standard Model and General Relativity?*
- ▶ **String theory** seems to reproduce both simultaneously.
- ▶ Starting in 1960's, string theory produced bosons in  $D = 26$ .
- ▶ In 1980's, super(symmetric)string theory also produced fermions in  $D = 10$ .
- ▶ In 1990's, J. Maldacena conjectured **AdS/CFT correspondence**.

**Anti-de Sitter spacetime**  $\leftrightarrow$  **Conformal field theory**  
Type IIB  $\text{AdS}_5$  superstring  $\quad \mathcal{N} = 4$  Super-Yang-Mills

- ▶ In this talk, review [AF09] approach to quantising  $\text{AdS}_5 \times S^5$  superstring.

$$\mathcal{L} \longrightarrow \mathcal{H} \longrightarrow \hat{\mathcal{H}} \longrightarrow \dots$$


# Why $\text{AdS}_5 \times S^5$ ?



**Figure 1.** Classical strings on hypersurfaces  $\text{AdS}_n$  and  $S^n$  for  $n = 2$  [Tse11, Sok16].

# Bosonic string theory

- ▶ Generalise relativistic point particle to one-dimensional string.



**Figure 2.** Worldline of particle and worldsheet of a closed string.

- ▶ Parameterise spacetime coordinates  $X^M(\sigma)$  where  $\sigma = (\tau, \sigma)$ ,  $M = 1, \dots, 26$  and

$$\partial_\tau X^M = \dot{X}^M, \quad \partial_\sigma X^M = X'^M.$$

- ▶  $G_{MN} \sim$  spacetime metric and  $\gamma_{\alpha\beta} \sim$  metric on the **worldsheet** for  $\alpha, \beta \in \{\tau, \sigma\}$ .
- ▶ For closed strings, worldsheet parameterised by  $\tau \in \mathbb{R}$  and  $-\pi r < \sigma < \pi r$ .

- ▶ **Polyakov action** for bosonic strings [GSW88]:

$$S = \int d\tau \int d\sigma \mathcal{L} = -\frac{T}{2} \int d\tau d\sigma \gamma^{\alpha\beta} \partial_\alpha X^M \partial_\beta X^N G_{MN}.$$

- ▶ Solving equations of motion  $\delta S / \delta \gamma_{\alpha\beta} = 0$  yields **Virasoro constraints**

$$C_1 = p_M X'^M = 0, \quad C_2 = p_M p^M + T^2 X'_M X'^M = 0.$$

- ▶ Rewrite Polyakov action in **first-order form** as

$$\begin{aligned} S &= \int d\tau d\sigma \left( p_M \dot{X}^M + \frac{\gamma^{\tau\sigma}}{\gamma^{\tau\tau}} C_1 + \frac{1}{2T\gamma^{\tau\tau}} C_2 \right) \\ &= \int d\tau d\sigma p_M \dot{X}^M. \end{aligned}$$

# Superstring theory

Bosonic/fermionic encoded in **superalgebra** structure [Kac77].

- $\mathcal{G} = \mathcal{G}^{(0)} \oplus \mathcal{G}^{(1)}$  is  $\mathbb{Z}_2$ -graded such that

$$[\mathcal{G}^{(a)}, \mathcal{G}^{(b)}] \subseteq \mathcal{G}^{(a+b)} \quad \text{mod } \mathbb{Z}_2 = \{\mathbf{0}, \mathbf{1}\}.$$

- If  $M \in \mathcal{G} \equiv \mathfrak{su}(2, 2|4)$ , then  $MH + HM^\dagger = 0$  and  $\text{str}(M) = \text{tr}(m) - \text{tr}(n) = 0$  where

$$M = \begin{pmatrix} \text{even} & \text{odd} \\ m & \theta \\ \eta & n \end{pmatrix}, \quad H = \begin{pmatrix} \gamma^5 & 0 \\ 0 & \mathbb{1}_4 \end{pmatrix}.$$

- Refine to  $\mathbb{Z}_4$ -grading with decomposition

$$\mathcal{G} = \mathcal{G}^{(0)} \oplus \mathcal{G}^{(1)} \oplus \mathcal{G}^{(2)} \oplus \mathcal{G}^{(3)},$$

$$M = M^{(0)} + M^{(1)} + M^{(2)} + M^{(3)},$$

according to automorphism  $\Omega : \mathcal{G} \rightarrow \mathcal{G}$  where  $\Omega^4(M) = M$  and

$$M^{(k)} = \frac{1}{4} \left[ M + i^{3k} \Omega(M) + i^{2k} \Omega^2(M) + i^k \Omega^3(M) \right].$$

# Superstring theory

- ▶ Parameterise  $\text{AdS}_5 = \{t, z^i\}$  and  $S^5 = \{\phi, y^i\}$  for  $i = 1, \dots, 4$ .
- ▶ Collect  $X^M \in \{t, \phi, x^\mu\}$  where  $x^\mu \sim$  transversal degrees of freedom for  $\mu = 1, \dots, 8$ .
- ▶ Define current  $A_\alpha = -\mathfrak{g}^{-1} \partial_\alpha \mathfrak{g} \in \mathcal{G}$  where  $\mathfrak{g} = \mathfrak{g}_b(t, \phi, x^\mu) \mathfrak{g}_f(\chi) \in G = \exp \mathcal{G}$ .

**Green-Schwarz action** for  $\text{AdS}_5 \times S^5$  superstring

$$S = -\frac{T}{2} \int d^2\sigma \left[ \gamma^{\alpha\beta} \text{str}(A_\alpha^{(2)} A_\beta^{(2)}) + \kappa \varepsilon^{\alpha\beta} \text{str}(A_\alpha^{(1)} A_\beta^{(3)}) \right]$$

- ▶ Introduce auxiliary  $\pi = \pi^{(2)} \in \mathcal{G}^{(2)}$ , the first-order form is

$$S = \int d^2\sigma \left[ -\text{str}(\pi A_\tau^{(2)}) + \frac{\gamma^{\tau\sigma}}{\gamma^{\tau\tau}} C_1 + \frac{1}{2T \gamma^{\tau\tau}} C_2 \right] - \frac{T}{2} \int d^2\sigma \kappa \varepsilon^{\alpha\beta} A_\alpha^{(1)} A_\beta^{(3)}$$

where this time

$$C_1 = -\text{str}(\pi A_\sigma^{(2)}) = 0, \quad C_2 = \text{str}(\pi^2 + T^2 A_\sigma^{(2)} A_\sigma^{(2)}) = 0.$$

# Gauge fixing

	$\kappa$ -symmetry gauge ( <b>s</b> )	Light cone gauge ( <b>s, b</b> )
Gauge freedom	$\delta\mathcal{L} = 0$ under $\mathfrak{g} \rightarrow \mathfrak{g}e^{\epsilon(\tau,\sigma)}$ provided $\kappa = \pm 1$	$\delta\mathcal{L} = 0$ under $(\tau, \sigma) \rightarrow (\tilde{\tau}, \tilde{\sigma})$ diffeomorphisms
Gauge fixing	$\chi \rightarrow \left( \begin{array}{cc cc} 0 & 0 & 0 & a \\ 0 & 0 & b & 0 \\ \hline 0 & b^\dagger & 0 & 0 \\ -a^\dagger & 0 & 0 & 0 \end{array} \right)$	$x_+ = \frac{1}{2}(\phi + t) \rightarrow \tau,$ $p_+ = \frac{1}{2}(p_\phi - p_t) \rightarrow 1.$



# Gauge fixing

- ▶ Recall for bosonic string, first-order form of action is

$$\begin{aligned} S &= \int d\tau d\sigma p_M \dot{X}^M = \int d\tau d\sigma (p_\mu \dot{x}^\mu + p_t \dot{t} + p_\phi \dot{\phi}) \\ &= \int d^2\sigma (p_\mu \dot{x}^\mu + p_- \dot{x}_+ + p_+ \dot{x}_-). \end{aligned}$$

In light cone gauge, where  $x_+ = \tau$  and  $p_+ = 1$ ,

$$S = \int d^2\sigma (p_\mu \dot{x}^\mu + p_- + \cancel{\dot{x}_-}) = \int d^2\sigma (p_\mu \dot{x}^\mu - \mathcal{H}).$$

- ▶ Hamiltonian of classical bosonic string in light cone gauge is

$$\mathcal{H} = -p_-(p_\mu, x^\mu, x'^\mu).$$

- ▶ What is  $p_-$  for superstring?

# Perturbative quantisation

- ▶ For superstring, one finds

$$\mathcal{H} = \dots = \frac{i}{2} \text{str}(\pi \Sigma_+ \mathfrak{g}(x^\mu) (\mathbb{1} + 2\chi^2) \mathfrak{g}(x^\mu))$$

↑  
3 weeks

$$+ \kappa \frac{T}{2} (G_+^2 - G_-^2) \text{str}(\Sigma_+ \chi \sqrt{\mathbb{1} + \chi^2} \mathcal{K} F_\sigma^{st} \mathcal{K}^{-1})$$
$$- \kappa \frac{T}{2} G_\mu G_\nu \text{str}(\Sigma_\nu \Sigma_+ \chi \sqrt{\mathbb{1} + \chi^2} \Sigma_\mu \mathcal{K} F_\sigma^{st} \mathcal{K}^{-1}).$$

- ▶ Not nice! Trick: rescale  $\sigma \rightarrow \sigma T$  such that  $2\pi r \rightarrow 2\pi r T$  and also rescale

$$x^\mu \rightarrow x^\mu / \sqrt{T}, \quad p_\mu \rightarrow p_\mu / \sqrt{T}, \quad \chi \rightarrow \chi / \sqrt{T}.$$

- ▶ End up with perturbative expansion in large tension limit

$$S = \int d^2\sigma \left( \mathcal{L}_2 + \frac{1}{T} \mathcal{L}_4 + \frac{1}{T^2} \mathcal{L}_6 + \mathcal{O}(T^{-3}) \right) \approx \int d^2\sigma \mathcal{L}_2.$$

# Perturbative quantisation

- Part of Lagrangian quadratic in original fields  $(x^\mu, p_\mu, \chi)$  is

$$\mathcal{L}_2 = p_\mu \dot{x}^\mu - \frac{i}{2} \text{str}(\Sigma_+ \chi \dot{\chi}) - \mathcal{H}_2,$$

where we read off

$$\mathcal{H}_2 = \left[ \frac{1}{4} P_{a\dot{a}} P^{a\dot{a}} + Y_{a\dot{a}} Y^{a\dot{a}} + Y'_{a\dot{a}} Y'^{a\dot{a}} + \frac{1}{4} P_{\alpha\dot{\alpha}} P^{\alpha\dot{\alpha}} + Z_{\alpha\dot{\alpha}} Z^{\alpha\dot{\alpha}} + Z'_{\alpha\dot{\alpha}} Z'^{\alpha\dot{\alpha}} \right] \\ + \left[ \eta_{\alpha\dot{\alpha}}^\dagger \eta^{\alpha\dot{\alpha}} + \frac{\kappa}{2} \eta'_{\alpha\dot{\alpha}} \eta'^{\alpha\dot{\alpha}} - \frac{\kappa}{2} \eta_{\alpha\dot{\alpha}}^\dagger \eta'^{\dagger\alpha\dot{\alpha}} + \theta_{a\dot{\alpha}}^\dagger \theta^{a\dot{\alpha}} + \frac{\kappa}{2} \theta'_{a\dot{\alpha}} \theta'^{a\dot{\alpha}} - \frac{\kappa}{2} \theta_{a\dot{\alpha}}^\dagger \theta'^{\dagger a\dot{\alpha}} \right].$$

8 Bosons

- Can now promote fields to operators, i.e. **quantisation** of the superstring:


$$Y^{a\dot{a}}(\tau, \sigma) = \int_{\mathbb{R}} \frac{dp}{2\pi} \frac{1}{\sqrt{2E_p}} \left( e^{i\sigma p} a_p^{a\dot{a}} + e^{-i\sigma p} \varepsilon^{ab} \varepsilon^{\dot{a}\dot{b}} a_{b\dot{b},p}^\dagger \right),$$

$$\theta^{a\dot{\alpha}}(\tau, \sigma) = \int_{\mathbb{R}} \frac{dp}{2\pi} \frac{1}{\sqrt{2E_p}} \left( e^{i\sigma p} f(p) a_p^{a\dot{\alpha}} + e^{-i\sigma p} h(p) \varepsilon^{ab} \varepsilon^{\dot{\alpha}\dot{\beta}} a_{b\dot{\beta},p}^\dagger \right).$$

8 Fermions

# Perturbative quantisation

- ▶ In decompactification limit ( $T \rightarrow \infty$ ) the worldsheet cylinder becomes a plane.



**Figure 3.** String excitation in decompactification limit

- ▶ Nice! Reduced problem to 1+1-dimensional QFT with 8 bosons and 8 fermions.
- ▶ What about **full** spectrum?

## AdS/CFT

$$\mathcal{L} \longrightarrow \mathcal{H} \longrightarrow \hat{\mathcal{H}} \longrightarrow \text{Spectrum} \leftrightarrow \mathcal{N} = 4 \text{ SYM}$$

First-order formalism

$\kappa$ -symmetry gauge

Light cone gauge

Thermodynamic Bethe  
Ansatz (TBA)

Perturbative quantisation  
in decompactification limit



Thank you

# References

- [AF09] Gleb Arutyunov and Sergey Frolov. *Foundations of the  $AdS_5 \times S^5$  Superstring. Part I*. 2009. arXiv: [0901.4937 \[hep-th\]](#).
- [Tse11] A. A. Tseytlin. *Review of AdS/CFT Integrability, Chapter II.1: Classical  $AdS_5 \times S^5$  string solutions*. 2011. arXiv: [1012.3986 \[hep-th\]](#).
- [Sok16] Leszek M. Sokolowski. *The bizarre anti-de Sitter spacetime*. 2016. arXiv: [1611.01118 \[gr-qc\]](#).
- [GSW88] M.B. Green, J.H. Schwarz, and E. Witten. *Superstring Theory: Volume 1, Introduction*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1988. ISBN: 9780521357524.
- [Kac77] V.G Kac. “Lie superalgebras”. In: *Advances in Mathematics* 26.1 (1977).

- Where is the spacetime hiding?

$$\text{AdS}_5 \times S^5 \cong \frac{SO(4,2)}{SO(4,1)} \times \frac{SO(6)}{SO(5)} \cong \frac{SU(2,2)}{SO(4,1)} \times \frac{SU(4)}{SO(5)} \subset \frac{SU(2,2|4)}{SO(4,1) \times SO(5)}.$$

- Automorphism  $\Omega(M) = -\mathcal{K}M^{st}\mathcal{K}^{-1}$  where

$$M^{st} = \begin{pmatrix} m^t & -\eta^t \\ \theta^t & n^t \end{pmatrix}, \quad \text{and} \quad \mathcal{K} = \begin{pmatrix} K & 0 \\ 0 & K \end{pmatrix}, \quad K = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

- Decomposition

$$M^{(0)} = \frac{1}{2} \begin{pmatrix} m - Km^tK^{-1} & 0 \\ 0 & n - Kn^tK^{-1} \end{pmatrix}, \quad M^{(1)} = \frac{1}{2} \begin{pmatrix} 0 & \theta - iK\eta^tK^{-1} \\ \theta + iK\eta^tK^{-1} & 0 \end{pmatrix},$$

$$M^{(2)} = \frac{1}{2} \begin{pmatrix} m + Km^tK^{-1} & 0 \\ 0 & n + Kn^tK^{-1} \end{pmatrix}, \quad M^{(3)} = \frac{1}{2} \begin{pmatrix} 0 & \theta + iK\eta^tK^{-1} \\ \theta - iK\eta^tK^{-1} & 0 \end{pmatrix}.$$



# Appendix

- Embedding coordinates. Set  $\mathfrak{g}_b(t, \phi, x^\mu) = \Lambda(t, \phi)\mathfrak{g}(x^\mu)$  where

$$\Lambda(t, \phi) = \exp \frac{i}{2} \begin{pmatrix} t\gamma^5 & 0 \\ 0 & \phi\gamma^5 \end{pmatrix}, \quad \mathfrak{g}(x^\mu) = \begin{pmatrix} \frac{1}{\sqrt{1-\mathbf{z}^2/4}}[\mathbb{1}_4 + \frac{1}{2}z^i\gamma^i] & 0 \\ 0 & \frac{1}{\sqrt{1+\mathbf{y}^2/4}}[\mathbb{1}_4 + \frac{1}{2}y^i\gamma^i] \end{pmatrix},$$
$$\mathfrak{g}_f(\chi) = \chi + \sqrt{\mathbb{1}_8 + \chi^2}.$$

- Light cone coordinates

$$\begin{aligned} t &= x_+ - ax_-, & x_+ &= a\phi + (1-a)t, \\ \phi &= x_+ + (1-a)x_-, & x_- &= \phi - t. \end{aligned}$$

- Isomorphism

$$\mathfrak{su}(4) \sim \mathfrak{so}(6) = \text{span}_{\mathbb{R}} \left\{ \frac{i}{2}\gamma^i, \frac{1}{4}[\gamma^i, \gamma^j] \right\}, \quad i, j = 1, \dots, 5,$$
$$\mathfrak{su}(2, 2) \sim \mathfrak{so}(4, 2) = \text{span}_{\mathbb{R}} \left\{ \frac{1}{2}\gamma^i, \frac{i}{2}\gamma^5, \frac{1}{4}[\gamma^i, \gamma^j], \frac{i}{4}[\gamma^i, \gamma^5] \right\}, \quad i, j = 1, \dots, 4.$$